

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Developmental Biology

journal homepage: www.elsevier.com/locate/developmentalbiology

Corrigendum

Corrigendum to: “Six1 transcription factor is critical for coordination of epithelial, mesenchymal and vascular morphogenesis in the mammalian lung” [Dev. Biol. 353 (2011) 242–258]



Ahmed H.K. El-Hashash^{a,*}, Denise Al Alam^a, Gianluca Turcatel^a, Orquidea Rogers^a,
Xue Li^{b,c}, Saverio Bellusci^a, David Warburton^{a,**}

^a Developmental Biology and Regenerative Medicine Program, Saban Research Institute, Children's Hospital Los Angeles, Keck School of Medicine of University of Southern California, 4650 Sunset Boulevard MS35, Los Angeles, CA 90027, USA

^b Department of Urology, Children's Hospital Boston, and Department of Surgery and Pathology, Harvard Medical School, Boston, MA, USA

^c Harvard Stem Cell Institute, Harvard University, Cambridge, MA, USA

Figs. 6 and 9 were incorrect as printed. For the reader's convenience, the correct Figs. 6 and 9 are printed here.




Fig. 6. Reduced expression of distal epithelial progenitor markers in Six1 (–/–) lungs. (A, B, E, F, I and J) Immunohistochemistry shows marked reduction of the expression levels of epithelial progenitor markers Sox9, Id2 and N-myc at distal epithelial tips of E14.5 Six1 (–/–) lungs (B, F and J; arrows), compared to control lungs (A, E and I; arrows). This decrease was also demonstrated at gene levels by real-time PCR for Sox9 (K). (C, D, G and H) Immuno-cytochemistry shows increased number of cells that express Sox9 (D) and Id2 (H) after Six1 overexpression in MLE-15 cells for three days in culture. Blue nuclear labeling with DAPI in C, D, G and H. (L) Quantification of experiments shown in C and D for Sox9 and for PHH-3-positive mitotic MLE15 cells. *Significantly different from control ($n=527$; $P < 0.05$; Student's t -test). Scale bars: 50 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

DOI of original article: <http://dx.doi.org/10.1016/j.ydbio.2011.02.031>

* Corresponding author.

** Correspondence to: Developmental Biology, Regenerative Medicine and Surgery Program, Saban Research Institute, Children's Hospital Los Angeles, Keck School of Medicine and School of Dentistry, 4661 Sunset Boulevard MS 35, Los Angeles, CA 90027, USA. Fax: +1 323 361 3613.

E-mail addresses: aelhashash@chla.usc.edu (A.H.K. El-Hashash), dwarburton@chla.usc.edu (D. Warburton).

<http://dx.doi.org/10.1016/j.ydbio.2014.04.009>

0012-1606/© 2014 Elsevier Inc. All rights reserved.




Fig. 9. Defective smooth muscle integrity in *Six1*($-/-$) embryonic lungs. (A–H) Immunohistochemistry with α -SMA antibody shows severely reduced staining surrounding blood vessels (B; black arrows) and large bronchial airways (F; black arrows) in *Six1*($-/-$) lungs compared to control lungs (A and E; black arrows). Note several breaches in the blood vessel walls, (B and D; blue arrows) and thin smooth muscle layer with herniation of the intact endothelial cell layer (B; yellow arrows) in mutant lungs. (C, D, G and H) α -SMA/Haematoxylin staining shows that spindle-shaped smooth muscle cells present in *Six1*($-/-$) lungs surrounding blood vessels and large bronchia (D and H; black arrows) compared to control lungs (C and G; black arrows). Note that sections are weakly stained with α -SMA antibody in order to see cell morphology in C and G. (I–J) Staining with PECAM antibody shows comparable signals in wildtype and *Six1*($-/-$) lungs (I and J; black arrows). Scale bars: 50 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)